

Minireview Article

Intestinal Invaginations in Adults

Abstract

Intestinal invaginations in adults are rare phenomena. The most important etiological causes are malignancies. The most common symptoms are abdominal pain, nausea, vomiting, inability to pass flatus and defecation, and distention in the abdomen due to intestinal obstruction. On physical examination, abdominal distention, decreased bowel sounds, and sometimes a mass may be palpable on deep palpation.

In direct radiography; air-liquid levels can be seen. A mass can be detected in the invagination area on ultrasonography. Definitive diagnosis is usually made by computerized tomography.

Treatment of intestinal invaginations in adults is usually surgery. It can be done by laparoscopic or open surgery.

As a result, we think that when a definitive diagnosis is made in adults, intestinal invagination should be treated surgically in accordance with oncological principles, since the etiological factor is mostly malignancy.

Keywords: Iliocecal, adult invagination, intussusception, diagnosis, treatment.

Introduction

Intestinal invagination (intussusception) is intertwinement of the intestinal segments like a telescope. Invaginations may occur in ileoileal, ileocecal or colocolic form. The most common type is ileocecal invaginations.¹⁻⁴ Intestinal intussusception are especially common in children under 1 year of age.⁵ They are rare in adults and are about 1-5% cause of bowel obstruction.^{1,3,4}

A wide variety of factors might play a role in intestinal invaginations; The most important cause of intestinal invaginations in adults is malignancies.^{1,3,4,6-12} It has also been reported that the etiologic cause is malignancy, especially in colonic invaginations.¹³

In a study by Cherni et al., it was reported that nitric oxide (NO) is the major inhibitory neurotransmitter in the enteric nervous system, inflammatory events cause excessive NO release in cases of ileocecal invagination, which results in severe relaxation of the ileocecal valve and ultimately ileocecal invaginations.¹⁴ In a similar study by Kaemmerer et al., it was reported that adenovirus infections cause CD3 positive lymphocyte infiltration, which causes enlargement of peyer's patches and mesenteric lymph nodes due to inflammatory neuropathy in the mesenteric plexus, which is the cause of ileocecal invaginations.⁵

There are cases that are etiologically less common in adults, including intraluminal lipoma in the ileum¹⁵⁻¹⁷, endometriosis in the appendix and ileum¹⁸, inflammatory fibroid polyp (Vanek tumor)^{19,20}, due to colonoscopy²¹, Meckel's diverticulum²², intestinal tuberculosis⁴, transmural diffuse large B cell lymphoma infiltration²³, idiopathic²⁴, submucosal colonic lipoma^{2,25}, Chron's disease²⁶, leiomyoma in the ileum²⁷, Burkitt lymphoma²⁸, cases of induced intestinal invagination have been reported.

A case of double ilio-caeco-colic invagination due to tumor was reported in a case report by Saclo et al.²⁹

Zhang et al. reported a very rare case of duodeno-duodenal invagination in a 23-year-old male patient.³⁰

In cases of invagination, patients usually admitted to the hospital with symptoms such as abdominal pain, nausea, vomiting, inability to pass flatus and defecation, and abdominal distention due to intestinal obstruction.

Depending on the obstruction site in the intestines, air-fluid levels may appear on the abdomen x-ray. Enlarged bowel loops and air-fluid levels may be revealed on ultrasonography.^{1,6}

The “pseudo- kidney” appearance formed by the invaginated intestines on abdominal ultrasonography might be diagnostic.⁷

In intestinal invaginations, CT has the high sensitivity and specificity rates in diagnosis.^{2,6,15,17,19,28}

According to the results of the study by Peters et al., CT alone is sufficient for a definitive diagnosis in intestinal invaginations.¹⁵

According to a study by Treppiedi et al., using "contrast-enhanced tomography" instead of classical CT in intestinal invaginations can significantly increase the rate of correct diagnosis.³¹

Although ileocecal invaginations are rare in adults, they are cases that should be handled carefully during follow-up and treatment, as malignant causes usually play a role in etiology.

Therefore, some authors do not recommend attempting a reduction before or during the operation considering the oncological risks.^{3,7,8}

However, in a retrospective study Honjo et al. reported that 28 of 44 adult invagination cases had reduction, and reduction should be attempted before or during surgery.¹

Yakan et al. reported that reduction should only be attempted in ilioileal intussusceptions without a suspected malignancy in their study in an invagination case series of 20 cases.¹³

In cases of invagination, reduction is generally performed by pneumatic or enema.^{31,32}

Treppiedi et al. reported in their study that pneumatic reduction should be tried before surgery in cases of invagination.³¹

Zhang et al. reported that they achieved successful results with endoscopic reduction in a case of duodeno-duodenal invagination.³⁰ Surgical applications in intestinal invaginations may differ according to the etiological factor, open or laparoscopic surgery can be applied.³²

The width of the operation to be performed should be kept large considering the risk of malignancy, and "en bloc" resection should be performed.^{2,8,29}

Honjo et al. in 12 adult intussusception cases¹, Saito et al. in a case caused by an intestinal pedicled lipoma¹⁶, Hong performed laparoscopic resection and anastomosis in a case of lipoma-induced invagination.¹⁷ However, as seen in the literature reviews, there are no cases of laparoscopic iliocecal invagination resection in large series in adults yet.

In a study conducted by Bahman, reduction was tried 3 times with an interval of 2-4 hours in invagination cases, successful results were obtained in 85% of the cases, and 52 cases of unsuccessful intestinal invagination were randomly divided into 2 groups: Laparoscopic surgery was performed in the first group of 26 people, and open surgery was performed in the other group of 26 people, and the duration of the operation, the transition period to oral nutrition, hospitalization times, wound infection rates were compared. According to the results obtained in the study; operation times were found to be significantly higher in the laparoscopic group compared to open surgery, and no significant difference was found between the two groups in terms of other criteria.³² For this reason, it has been reported

that reduction should be attempted primarily in cases of intestinal invagination and that laparoscopic surgery is as safe as open surgery.

Conclusion

According to the results of our study, the most common cause of intestinal invaginations in adults is malignancies. CT is the gold standard in diagnosis. However, in many cases, the diagnosis can be made with USG. In invagination cases with suspected malignancy, oncological principles should be followed. Open or endoscopic methods can be used safely in surgical treatment.

References

1. Honjo H, Hiroshi M, Nobuyasu K: Adult Intussusception: A Retrospective Review. *World Journal of Surgery*.2015;39(1):134-138.
2. Cordeiroa J, Corderioa L, Possaa P, Candidoa P, Oliveirab A: Intestinal intussusception related to colonic pedunculated lipoma: A case report and review of the literature. *International Journal of Surgery Case Reports*.2019; 55: 206-209.
3. Barut B, Yonder H, Sarıcı KB, Özdemir F, et al.: Retrospective analysis of cases of intestinal invagination treated and followed-up at our clinic .*North Clin Istanbul* 2017;4(2):141–144.

4. Kaval S, Singhai BM, Sanieev K, Chandra PS: Adult intussusception: An institutional experience and review of literature. *Archives of International Surgery*. 2014;4(1):25-30.
5. Kaemmerer E, Jens J, Tischendorf W, Steinau G, et al.: Ileocecal Intussusception with Histomorphological Features of Inflammatory Neuropathy in Adenovirus Infection. 2009;2009(2):1-4.
6. Lorenzo JG, Martinez AT, Ruiz L, Pastor F, et al.: Intestinal invagination in adults: preoperative diagnosis and management. *International Journal of Colorectal Disease*. 2004;19(1):68-72.
7. Verre L, Rossi R, Gaggelli I, Piccolomini A, Podzemny V, Trone A: Ileocecal-colonic intussusception caused by cecal adenocarcinoma A case report. *Ann Italiani di Chirurgia*, 2012; pii: S2239253X12019081.
8. Marion J, Pollheimer T, Baumgartner VS, Pollheimer JT, Lackner H: Ileocecal intussusception caused by lung cancer metastasis. *Wien Klin Wochenschrift*. 2009;121(11):413-416.
9. Carlos CS, Javier GS: Adult Intestinal Intussusception. *Diseases of the Colon & Rectum*. 2021;64(6):645-648.

10. Panaccio P, Fiordaliso M, Testa D, Mazzola L, Battilana M, Cotellese R, et al: Minimally Invasive Treatment of Sporadic Burkitt's Lymphoma Causing Ileocaecal Invagination. *Case Rep Surg*. 2018. <https://doi.org/10.1155/2018/6265182>.
11. Deena I, Nina PP, Patel P, Lotfipour S: Ileocecal Intussusception in the Adult Population: Case Series of Two Patients. *Western Journal of Emergency Medicine*. 2010; 11(2):197-200.
12. Gruenberger EH, Hassan M, Kratzke I: Long-tract ileocolic intussusception due to mucinous adenocarcinoma of the ileocecal valve: a case report and literature. 2017; 2017(3).
13. Yakan S, Çalışkan C, Makay O, Denecli AG, et al.: Intussusception in adults: Clinical characteristics, diagnosis and operative strategies. *World J Gastroenterol*. 2009; 15(16): 1985-1989.
14. Cherni T, Parana S, Puri P: New hypothesis on the pathogenesis of ileocecal intussusception. *Journal of Pediatric Surgery*. 2007; (42): 1515–1519.
15. Peters J: Ileocecal invagination-an unequivocal computer tomography diagnosis. *Digitale Bild diagnostik*. 1985; 5(4):195-196.

16. Saito K, Osawa, K Morohara, K Nakamura, et al.: Laparoscopy-Assisted Resection of Ileocecal Intussusception Caused by Ileal Pedunculated Lipoma. *International College of Surgeon*.2013;98(4):330-333.
17. Hong Yu, Shuodong W, Renzhong Z, Xiaopeng Y: Single-incision laparoscopic-assisted ileal resection for adult intussusception. *Journal of Minimal Access Surgery*.2016;12(2):182-185.
18. Dainius E, Pankratjevaite L, Bradulskis S, Venskutoniskis S: Intussusception of the Appendix and Ileum Endometriosis: A Case Report. *Visceral Medicine*.2016;32:211-213.
19. Chapa LAF, Salgado CLE, Garza CAA, Villafranca ARI, Alanís RÓC: Ileocecal intussusception in an adult patient caused by a Vanek tumor: Case report.2018;19(3).
20. Lasithiotakis, K, DG Evangelia, F Dimitrios, Athanasakis, I, Z Odysseas, Chalkiadakis, G: Ileocolic Intussusception Precipitated by Diagnostic Colonoscopy: A Case Report. *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques*.2012;22(3):161-163.
21. Amir W, Hassan W, Tech W: Intussusception after Colonoscopy: A Case Report and Review of Literature. *J of minimal Access surgery*.2018;51(6):591-595.

22. Morrison J, Jeanmonod R: Intussusception Secondary to a Meckel's Diverticulum in an Adolescent. *Case Reports on Emergency Medicine*. 2011; doi:10.1155/2011/623863
23. Saka R, Sasaki T, Matsuda I, Nose Satoko, et al.: Chronic ileocolic intussusception due to transmural infiltration of diffuse large B cell lymphoma in a 14-year-old boy: a case report. *AJE*. 2015;4:366.
24. Harma M, Harma M, Karadeniz G, Arıkan I, et al.: Idiopathic ileoileal invagination two days after cesarean section. *The Journal of Obstetrics and Gynecology Research*. 2011;37(2):160-162.
25. Ozen O, Guler Y, Yuksel Y. Giant colonic lipoma causing intussusception: CT scan and clinical findings. *Pan Afr Med J*. 2019;32(27).
26. Jabi R, Elmir S, Quryemchi M, Bouziane M: Acute Intestinal Invagination: An Exceptional Method of Revealing Crohn's Disease. *Cureus*. 2021;13(10).
27. Bouali M, Yousra EB, Bakouri AE, Bensardi F, Hattabi KE, Fadil A: Acute Intestinal Invagination in Grelic Leiomyoma. *Asian Journal of Research in Surgery*. 2021;5(2):14-17.

28. Panaccio P, Fiordaliso M, Testa D, Mazzola L, Battilana M, Cotellesse R, et al: Minimally Invasive Treatment of Sporadic Burkitt's Lymphoma Causing Ileocaecal Invagination. *Case Rep Surg*.2018. <https://doi.org/10.1155/2018/6265182>.
29. Sacko O, Sissoko M, Soumare L, Keita S, Koumare S, Konare B: Double Invagination Ileo Coeco Colic due to a Carcinoma of the Right Colon about a Case. *Surgical Science*.2020;11(11).
30. Zhang KS, Bansal J, Bansal A, Chitnavis V: Adult Intestinal Intussusception Caused by the Gastrojejunostomy Tube: An Endoscopically Treatable Phenomenon. *Case Reports in Gastrointestinal Medicine*.2021. <https://doi.org/10.1155/2021/4325443>
31. Treppiedi E, Cocchi L, Zimmitti G, Manzoni A, Seletti V, Bizzotto A, et al: Ileocolic invagination in adults: A totally minimally invasive endoscopic and laparoscopic staged approach. *J Minim Access Surg*.2020;16(1):87-89.
32. Bahman R: Comparision of Laparoscopic Surgery and Open Syrgery in Treatment of Invagination Needed for Surgery. KR-TBZMED Home School of Medicine Theses(M). Tabriz University Medical Sciences.2020.