

# Evaluation of Antinuclear Antibodies and Differential Leucocyte Count in ABO Typed Women with Spontaneous Abortion in Yenagoa, Bayelsa State

## ABSTRACT

**Aim:** This study ~~was aimed at~~ ~~to~~ ~~evaluating~~ the anti-nuclear antibodies, ~~and~~ differential leucocytes count in ABO typed women with spontaneous abortion (SA) in Yenagoa, Bayelsa State.

**Study design:** Cohort cross-sectional study.

**Place and Duration of Study:** Scanex Diagnostics No 58a Nikton Road, Kpansia. Yenagoa, Bayelsa State, between January, 2018 and August, 2018.

**Methodology:** A total of 150 females were recruited; 50 with unexplained history of SA (UHSA) and 50 with unexplained recent SA (URSA) formed the test groups while 50 apparently healthy females who have children, ~~without no any~~ recent or history of abortion were used as control. ~~5ml~~ ~~Five milliliters (5mls)~~ of venous blood was collected using standard ~~venepuncture~~ ~~venipuncture~~ technique from each participant, out of which ~~two milliliters (2mls)~~ was dispensed into ethylenediaminetetraacetic acid (EDTA) bottles, and was used for DLC by manual spread technique and ABO testing, using the tile method. The remaining 3mls of blood ~~in plain bottle~~ was used ~~for the analysis of to analyse~~ ANAs by ~~the~~ ELISA Assay technique.

**Results:** The study participants, UHSA, URSA and Control, showed ~~that the~~ age group of 31-35 years ~~to have~~ ~~the~~ majority of spontaneous abortion of 18 (36%), 13 (26%) and 12 (24%), respectively. In the Blood group study, majority ~~were of~~ blood group ~~were~~ B; 4 (40%), 3 (22.2%), 1 (16.6%) and O; 7 (22.5%), 3 (8.1%), 3 (9.6%), respectively. Out of the 9 blood group B, URSA subjects only 3 of them tested positive for ANAs, out of 10 blood group B UHSA subjects, only 4 of them tested positive for ANAs & ~~out~~ of the 6 blood group B control participants, only 1 of them tested positive for ANAs; ~~Out~~ of the 37 blood group O URSA subjects only 3 of them tested positive for ANAs, also out of the 31 blood group O UHSA subjects, only 7 tested positive for ANAs and out of the 6 blood group O control participants only 1 tested positive for ANAs. Out of the 50 subjects with URSA, only 6 tested positive for ANAs, also out of the 50 UHSA subjects, only 11 tested positive for ANAs and out of the 50 control participants, only 4 tested positive for ANAs. The mean ANAs value (0.13±0.11 and 0.20±0.03) in URSA and UHSA were significantly lower ( $p < 0.05$ ) ~~when~~ compared to the control (0.07±0.01) ( $p = 0.001$ ) subjects. Also, the mean monocyte value (3.12±1.70 and 2.54±1.71) in URSA and UHSA were significantly higher ( $p < 0.05$ ) when compared to the control (2.20±1.48) ( $p = 0.020$ ) subjects, respectively. There was no significant ( $p > 0.05$ ) difference between the mean neutrophil, lymphocytes and eosinophil values (54.56±10.25 and 56.28±11.12; 41.62±11.15 and 40.56±11.25; 0.32±0.94 and 0.62±1.01) in the same order when compared to ~~that of~~ the control (56.10±10.56; 41.08±10.58; 0.52±0.88) ( $p = 0.6750$ ;  $p = 0.8905$ ;  $p = 0.2736$ ) subjects in the same order.

**Conclusion:** ~~It is therefore concluded that age, ANAs, ABO blood group~~ Age, ANAs, ABO blood group, and monocyte could be responsible for unexplained SA. It is recommended that ANAs, Blood groups and DLC be ~~properly~~ ~~adequately~~ checked in early pregnancy, for this shall further aid the prediction and prevention of spontaneous abortion.

*Keywords: Antinuclear Antibodies, Differential Leucocyte Count, ABO, Women, Spontaneous Abortion, Yenagoa, Bayelsa State*

## 1. INTRODUCTION

Three or more consecutive pregnancy losses ~~is spontaneous abortion SA [1], many causes of SA is not are spontaneous abortion SA [1], many causes of SA are unknown~~. 1% of the couple are affected [2], some SA ~~are is~~ caused by anatomical conditions, chromosomal disorders, immune factors, ovarian factors, life-style factors and infection (ACOG, 2001), in connection with the later disease are risk of cardiovascular complications which can be treated in some couples [3], ~~though. However,~~ some never had a cause identified, and after extensive investigations, about 50-75% of cases of SA were unexplained [2]. Spontaneous abortion can also be clinically recognized as pregnancy loss before the 20th week of gestation [4]. Spontaneous abortion is also known as miscarriage, and it refers to a pregnancy that ends spontaneously before the foetus has reached a viable gestational age [5]. The World Health Organization (WHO) defines it as expulsion or extraction of an embryo or fetus weighing 500 g or less [6]. The term "foetus" will be used for most of this discussion, although the term "embryo" is the correct developmental term at  $\leq 10$  weeks of gestation. Analytical issues, risk factors and aetiology of spontaneous abortion can be reviewed and managed independently [6].

Many researches have been carried out to identify the underlying mechanisms of spontaneous abortion, and evidence reveals that an immunologic mechanism is involved in some miscarriages. However, several studies of ~~then~~ immune interactions at the foeto-maternal interface and genetic-epidemiologic studies documented an immunological background for many spontaneous abortion cases [7]. Several autoantibodies, produced by the immune system directed against one or more of the individual's ~~own~~ proteins have been investigated as possible influences on reproductive success and failure. Auto-antibodies may persist for many years in the circulation as a marker of a previous autoimmune attack, but their presence does not necessarily indicate a recent disease process. The anti-phospholipid antibodies such as lupus anticoagulant, anti-cardiolipin, antinuclear, anti-  $\beta 2$  glycoprotein are associated with spontaneous abortion or as possible factors involved in infertility [7]. Antinuclear antibodies (ANAs), also known as ~~an~~ antinuclear factor (ANF) [8], are autoantibodies that bind to ~~the~~ contents of the cell nucleus. ~~In normal individuals, the immune system produces antibodies to foreign proteins (antigens). The immune system produces antibodies to foreign proteins (antigens) in normal individuals~~ but not to human proteins (auto-antigens). In some individuals, antibodies to human antigens are produced [9].

Immunoglobulin G (IgG) is an antibody that is about 75% of antibodies in ~~the~~ serum of humans. It is ~~the~~ Mostly found in circulation [10]. The molecules of 1gG ~~is are~~ created and released by plasma B cells, ~~when~~ compared to immunoglobulin M (IgM), which is the largest antibody produced by vertebrates, and it appears in response to initial exposure to antigen [11]. 1gM are responsible for the reaction caused by incompatible blood transfusion, the clumping of red blood cells if the recipient of a blood transfusion receives blood that is not compatible with their blood type [12].

White blood cells (WBCs), are cells of the immune system, ~~;~~ they are also called leucocytes. Their role is to protect against infections and foreign ~~bodybodies~~. White blood cells are derived from the multipotent cells of the bone marrow. WBCs are found in the blood and the lymphatic system; they all have ~~a~~ nucleus, ~~which make them different from other blood cells which differentiating them from other blood cells that~~ does not have nuclei [13]. The amount of white blood cells in the blood ~~is a show of shows~~ a normal or diseased state, ~~;~~ it is known to be a ~~very importantn essential~~ aspect of complete blood count.

The physiological roles of ABO blood group antigens remain a mystery despite their clinical importance. People with the common blood type O express neither the A nor B antigen, and they are perfectly healthy. Many diseases have been linked to some ABO phenotypes. For example, the ABO phenotype linked with stomach ulcers (Blood group O) and gastric cancer

was linked with blood group A individuals. Another observation is that individuals with blood type O tend to have lower levels of the von Willebrand Factor (vWF), which is a protein involved in blood clotting ~~so these individuals—so these individuals~~ could suffer from haemorrhage [14]. Shortly after the ABO blood groups were discovered, attention was directed toward the possibility of harmful effects when mother and fetus have different blood. This is the only blood group in which antibodies are constantly, predictably, and obviously present in the serum of people who lack the antigen on the red blood cells (RBCs). Spontaneous abortion is a pregnancy that ends spontaneously before the fetus has gotten to a viable gestational age [5]. ABO incompatibility occurs in 20% of pregnancies, but only 20% ~~of these~~ develop the hemolytic disease, which is milder than Rh incompatibility that can lead to abortion in the uterus [15].

Spontaneous abortion (SA) is a common occurrence without predictive tools and diagnostic cause. Studies on arrays of haematological investigations that provide the necessary support to the primary activities are still not known, ~~but~~ However, in order to be able to know the pattern in diagnosing spontaneous abortion, previous studies have implicated some markers like antinuclear antibodies, differential ~~leucocyte-leukocyte~~ count and ABO blood groups, [16]. Therefore, this study was designed to ~~bring out~~ identify possible markers in predicting spontaneous abortion using antinuclear antibodies (IgG), ABO antigens and immune cells of interest.

## 2. MATERIALS AND METHODS

### 2.1 Study Design

This study is a convenience cohort ~~cross-cross~~-sectional ~~study-investigation~~ carried out among women with spontaneous abortion attending emergency clinics in Yenagoa, Bayelsa State, Nigeria. The participants that were recruited for this study were grouped into two test groups; women with recent spontaneous abortion and women with a history of spontaneous abortion. Apparently, healthy women who already have children with no history of spontaneous abortion constituted the control group.

### 2.2 Study Area

The study was carried out ~~at~~ the Federal Medical Centre Yenagoa, Bayelsa State, Nigeria. The hospital is the foremost tertiary health institution with referrals from all nooks and crannies of the State. A hospital with 425 bed complement and is blessed with hardworking and highly dedicated staff numbering 2,216 regular staff. Bayelsa state, is located within Latitude 4<sup>0</sup> 15<sup>1</sup> North and Latitude 5<sup>0</sup> and 23<sup>1</sup> South [17]. It is also within longitude 5<sup>0</sup> 22<sup>1</sup> West and 6<sup>0</sup> 45<sup>1</sup> East. It is bounded by Delta State on the East, Rivers State on the West and the Atlantic Ocean on the Southern parts. Bayelsa State has eight local government areas, viz: Yenagoa, Ekeremor, Brass, Nembe, Ogbia, Southern Ijaw, Sagbama, Kolokuma/Opokuma. According to the 2006 census figures, Bayelsa has a population of about 1.7 million people [17].

### 2.3 Study Population

The study population was made up of one hundred and fifty (150) participants within the age range of 21-60 years, consisting of 50 women with unexplained recent spontaneous abortion (URSA) and 50 women with an unexplained history of spontaneous abortion (UHSA) used as Test groups and 50 apparently healthy women who already have children with no history of spontaneous abortion as the control group. A well-structured questionnaire was used to obtain the demographic details of the participants. Informed consent was obtained from each participant before enrollment into this study.

### 2.4 Eligibility Criteria

#### 2.4.1 Inclusion Criteria

This study focused on subjects with an established history of spontaneous abortion more than two consecutive times and subjects who just had spontaneous abortion (unexplained history and recent spontaneous abortion). The control group were apparently healthy females who already have had children without any history of spontaneous abortion.

#### 2.4.2 Exclusion Criteria

Exclusion criteria include the following: subjects with history of surgeries, subjects receiving radiotherapy, subjects under long term local and systemic drug therapy, HIV- positive individuals, subjects with systemic illness, subject with failing organs, subjects that was-were positive to urine glucose.

#### 2.5 Sample Size Calculation

The sample size for the study was determined using the formula:

$$N = \frac{(Z^2 Pq)}{d^2} \quad [18]$$

Where:

N = Minimum number of subjects

Z = Standard Normal deviation at 95% confidence interval usually set at 1.96

q = 1 - p

d = Confidence level 0.05

P = Prevalence of spontaneous abortion incidence in Nigeria is 100.0 per 100,000 abortion cases

Therefore  $\frac{100}{100,000} \times 100$

=  $\frac{10}{100} = 0.10$

P = 0.10

q = 1 - 0.10

$N = \frac{(1.96)^2 \times 0.10 \times (1-0.10)}{(0.05)^2}$

N = 138.3

= 138.

The sample size was supposed to be 138 participants, but due to the difficulty in getting test subjects, a convenience sample size of 100 was used for the test group.

#### 2.6 Sample Collection

Five millilitres (5mls) of blood was collected from each participant using the standard venepuncture technique as described by [6], with a sterile disposable needle. Three millilitres (3mls) of blood was dispensed into plain bottles and, the blood in plain bottles was allowed to clot to generate serum, the serum was separated by a Pasteur pipette Pasteur pipette separated the serum for antinuclear antibodies. The separated serum was centrifuged in the laboratory using a bench centrifuge at 4000rpm for two minutes to harvest clear serum. The bottles were labelled appropriately for each participant and stored in a freezer/cold store at -15°C to -30°C (- 20°C laboratory freezer) until analyzed. Prior to use, the samples were thawed quickly The samples were thawed quickly before use, preferably to 8°C and no warmer than room temperature. Also, 2mls of whole blood was dispensed into ethylene diamine tetraacetic acid (EDTA) bottles (di-potassium salts of EDTA, the optimal concentration is 1.5mg per ml of whole blood) for differential white blood cell count.

#### 2.7 Laboratory Analysis

##### 2.7.1 Determination of Antinuclear Antibodies

**Method:** Diagnostic Automation, Incorporation ANAs screen Enzyme-linked Immunosorbent Assay (ELISA) test system designed to detect IgG class antibodies was employed.

**Principle:** Diluted human serum was added to wells coated with purified nuclear antigens. Antinuclear Antibodies IgG specific antibody, if present bound to the antigen. All unbound materials were washed away, and the enzyme conjugate was added to bind to the antibody antibody-antigen complex, if present. Excess enzyme conjugate was washed off, and the

substrate was added. The plate was incubated to allow hydrolysis of the substrate by the enzyme. The intensity of the colour is directly proportional to the amount of IgG specific [antibody-antibodies](#) in the sample.

### **2.7.2 ABO/ Rhesus Testing**

**Method:** Tile Method

**Principles:**

The ABO grouping system is based on haemagglutination reaction. When red blood cells carrying antigens are exposed to the corresponding antibodies, they interact with each other to form a visible agglutination or clumping.

### **2.7.3 Methodology for Differential Leucocyte Count**

**Method:** Manual Spread

**Principles:** The manual differential [leucocyte-leukocyte](#) count is performed to determine the relative number of each type of leucocytes present in the blood. Studies of red blood cells, white blood cells and platelet morphology ~~is-are~~ also performed. The combination of polychrome methylene blue and eosin stains has selective staining properties. The differential staining allows one to identify the types of leucocytes on the smear.

## **2.8 Data Analysis**

~~Questionnaire~~ A questionnaire was used to gather information from the participants. Analysis of the data obtained in the study was done using the Statistical Package for the Social Sciences (SPSS) version 22.0. Student's t-test, Analysis of variance (ANOVA), Tukey's post hoc, and Pearson Correlation tests were used to test the significance of the data. Results were presented as mean  $\pm$  standard deviation and summarized as frequency (percentage). A probability value  $p < .05$  was considered statistically significant.

## **3. RESULTS AND DISCUSSION**

Antinuclear antibodies were initially identified as ~~a marker~~ markers to various disorders like auto-immune hepatitis, systemic lupus ~~erythematosus-erythematosus~~ and sharp syndrome. However, spontaneous abortion is now known to occur when there is an underlying auto immune process that affects the development of the placenta. The association of antinuclear antibodies with spontaneous abortion has been studied by several researchers using various research methods. However, there has not been a clear cut acceptable method in tackling this problem-. This probably ~~have-has~~ led to various data obtained by different studies. This work was carefully designed to study the incidence of antinuclear antibodies, and differential leucocyte count in ABO typed women with spontaneous abortion, women with unexplained recent spontaneous abortion (URSA), unexplained history of spontaneous abortion (UHSA) and their values compared with apparently healthy women, who already have children with no history of abortion (control).

**Table 1a: Demographic Characteristics of Study Participants**

Characteristics	Overall N (%)	CONTR OL N = 50 N( % )	URSA N = 50 N( % )	UHSA N = 50 N( % )
-----------------	------------------	---------------------------------	--------------------------	--------------------------

<b>Age (Mean ± SD)</b>	150 (100)	50 (33.3)	50 (33.3)	50 (33.3)
<b>Age Range</b>	33.89±7.61	35.58±8.97	33.14±7.24	32.95±6.63
21 – 25			9(18)	10(20)
26 – 30	24(12.6)	5(10)	10(20)	8(16)
31 – 35	28(18.6)	10(2)	13(26)	18(36)
36 – 40	43(28.6)	12(24)	8(16)	9(18)
41 – 45	24(16.0)	7(14)	7(14)	3(6)
46 – 50	15(10)	5(10)	2(4)	1(2)
51 – 55	10(6.6)	7(14)	1(2)	1(2)
56 – 60	5(3.3)	3(6)	0(0)	0(0)
<b>Marital Status</b>	1(2)	1(2)		
<b>Married</b>	150(100)	50(100)	50(100)	50(100)
<b>Educational Qualification</b>				
<b>Primary School</b>				
<b>Secondary School</b>	30(60)	8(26.6)	12(40)	10(33.3)
<b>Diploma</b>	37(74)	12(32.4)	13(35.1)	12(32.4)
<b>Degree</b>	24(48)	9(37.5)	7(29.1)	8(33.3)
<b>Others</b>	39(78)	10(25.6)	16(41.0)	13(33.3)
	20(40)	4(20)	6(30)	10(50)
<b>Ethnic Groups</b>				

*Note: Percentages may not add up to 100 due to rounding up; frequency for each variable may vary due to non-responses or missing values, % = Percentages, N = Number of study participants, URSA= unexplained recent spontaneous abortion, UHSA= unexplained history of spontaneous abortion.*

In this study, 12% were women with unexplained recent spontaneous abortion (URSA) recorded positivity for antinuclear antibodies, 22% were antinuclear antibodies positive in women with [unexplained history of spontaneous abortion \(UHSA\)](#), while only 8% of apparently healthy women were positive for antinuclear antibodies. This does not agree with the findings of Mohammed et al. [19], which established that the frequency of antinuclear antibodies was 31.8% in 110 women with history of spontaneous abortion and that only 5.7% of 35 healthy women with proven fertility and with no history of pregnancy loss or autoimmune disease, were positive. Many researchers believed that the presence of antinuclear antibodies acts as a risk factor for infertility, and antinuclear antibodies can be involved in the mechanisms causing [the failure of embryo implantation](#) [embryo implantation failure](#).

**Table 1b: Demographic Characteristics of Study Participants**

Characteristics	Overall N (%)	CONTR OL N = 50 N( % )	URSA N = 50 N( % )	UHSA N = 50 N( % )
Igbo	36(24)	0(0)	14(28)	22(44)
Hausa	7(4.6)	3(6)	4(8)	0(0)
Yoruba	17(11.3)	10(20)	2(4)	5(10)
Nembe	11(7.3)	6(34.3)	3(17.18)	2(11.4)
Epie	30(20)	6(12.6)	15(31.5)	9(18.9)
Ogbia	10(6.6)	5(31.5)	3(18.9)	2(12.6)
Ijaw	12(8)	5(26.2)	5(26.2)	2(10.5)
Ogoni	1(0.6)	0(0)	1(2)	0(0)
Kalabari	1(0.6)	0(0)	1(2)	0(0)
Ibibio	21(14)	15(30)	1(2)	5(10)
Isoko	4(2.6)	0(0)	1(2)	3(6)
Eshan	1(2)	0(0)	1(2)	0(0)
<b>Religion</b>				
Muslim	19(12.6)	8(16)	4(8)	7(14)
Christian	127(88)	42(84)	43(86)	42(84)
Others	4(2.6)	0(0)	3(6)	1(2)
<b>Occupation</b>				
Civil Servants	39(26)	16(41)	9(23)	20(51)
Housewife	5(3.3)	0(0)	2(4)	3(6)

Traders	22(14.6)	10(20)	5(10)	7(14)
Others	18(12)	10(20)	5(10)	3(6)

**Note:** Percentages may not add up to 100 due to rounding up; frequency for each variable may vary due to non-responses or missing values, % = Percentages, N = Number of study participants, URSA= unexplained recent spontaneous abortion, UHSA= unexplained history of spontaneous abortion.

On the assessment of the differential leucocyte count of individuals with unexplained spontaneous abortion and controls, the results of this study showed a higher significant difference ( $P=.02$ ) in the studied monocyte count in both unexplained recent (URSA) and history of spontaneous abortion (UHSA) when compared with the control (Table 2). This is in line with the work of Deng et al. [20], where they discovered that monocyte being the largest white blood cells, is involved in inflammation; they are more in chronic inflammation, viral infection and bacterial infection. Odutola et al. [21] also discovered that monocyte carries out phagocytic activities and they present antigen and produce cytokines. The clinical implication of low ANAs is found in chronic bacterial or viral infection and in patient with haematological malignancy haematological malignancy patients. It is a marker for spontaneous abortion; this is because inflammation of the uterus that does not allow it to be a suitable host for implantation of the embryo; natural killer cells mistake the fetal cells for cancer cells and attack them, making it un conducive for the fetus to thrive.

**Table 2: Comparison of Mean  $\pm$  Standard Deviation of ANAs, Differential Leukocyte Counts of Study Participants.**

Parameters/Unit	URSA (a)	UHSA (b)	Control(c)	F-value	p-value	TMC
ANAs(OD RATIO)	0.13 $\pm$ 0.11	0.20 $\pm$ 0.03	0.07 $\pm$ 0.01	7.747	0.001 (S)	a-c (S) b-c (S) a-b (S)
Neutrophils (%)	54.56 $\pm$ 10.25	56.28 $\pm$ 11.12	56.10 $\pm$ 10.56	0.3941	0.6750(NS)	ALL (NS)
Lymphocytes (%)	41.62 $\pm$ 11.15	40.56 $\pm$ 11.25	41.08 $\pm$ 10.58	0.1161	0.8905(NS)	ALL (NS)
Eosinophils (%)	0.32 $\pm$ 0.94	0.62 $\pm$ 1.01	0.52 $\pm$ 0.88	1.3080	0.2736(NS)	ALL (NS)
Monocyte (%)	3.12 $\pm$ 1.70	2.54 $\pm$ 1.71	2.20 $\pm$ 1.48	4.020	0.020 (S)	a-c (S) b-c (S) a-b (S)

**Key:** NS=Non Significant; S = Significant; TMC – Total Material Consumption (defines two types of data)  
ALL = As per every participant

In this study, there is a statistically higher significant difference ( $p=.02$ ) in antinuclear antibodies of all the participants as compared to that of the controls (Table 3). ANAs was positive in some spontaneous abortion individuals and also were positive in some spontaneous abortion individuals and control individuals, which may be due to some underlying immune disorders. This agrees with what Sarici et al. [22] reported, that ANAs is mainly associated with connective tissue diseases, and it is also found in healthy people most especially women and are mainly associated with connective tissue diseases, and it is also found in healthy people, especially women and the elderly. Kutteh and Odom [23] argued that spontaneous abortion reoccurs when the level of antinuclear antibodies is elevated in circulation, suggesting an autoimmune disorders, which causes an immune system to attack its own cells, tissue or organ by mistake. Antinuclear antibodies are not only present in patient with connective tissue disease but also present in a patient with connective tissue disease and in healthy people, according to Rai et al. [24].

**Table 3 Percentage Positivity & Negativity for Antinuclear Antibodies in the Study Population Irrespective of Blood Type.**

Parameters	No. of ANA Positivity	% of ANA Positivity	No. of ANAs Negativity	% of ANAs Negativity
URSA	6	12	44	88%
UHSA	11	22	39	78%

Control Subjects	4	8	46	92%
------------------	---	---	----	-----

**Key:** ANAs = Antinuclear Antibodies

~~It was also observed in this study~~ This study also observed that spontaneous abortion is highest in individuals within the age range of 31 – 35 years; 26% were of unexplained recent spontaneous abortion individuals, and 36% were of unexplained history of spontaneous abortion individuals (Table 1b). This is similar to the findings of Nyabo et al. [25]. It was also reported that spontaneous abortion risk increases with age, especially in women aged equal or greater than 35 years and that the frequency of its adverse effects in spontaneous abortion is about 10%.

In this study, hypersegmented neutrophil cells, acanthocytes, lymphocytes, schistocytes, polychromatic cells and macrocytic cells were common findings in women with spontaneous abortion (Table 2). This agrees with the study of Segolodi et al. [26], reported evidence of folate/vitamin B12 deficiency as a cause of spontaneous abortion. Hypersegmented neutrophils, macrocytes and macro ovalocytes are seen in macrocytic anaemia induced by vitamin B12 or folic acid deficiency. In the study of Yang et al. [27], it was reported that folic acid supplementation prevent megaloblastic anaemia during pregnancy and neural tube defects (NTDs) in the newborn infant. Thompson et al. [28] associated low Vitamin B12 with pregnancies affected by anencephaly (a defect in the formation of a baby's neural tube during development, baby born without parts of the brain and skull).

**Table 4 Percentage Positivity and Negativity for Antinuclear Antibodies in Blood Group B Subjects**

Parameters	No. of ANAs Positivity	% of ANAs Positivity	No. of ANAs Negativity	% of ANAs Negativity
URSA	3	22.2	47	94
UHSA	4	40	46	92
Control Subjects	1	16.6	49	98

**Key:** ANAs = Antinuclear Antibodies

According to Tamura and Picciano [29], folate deficiency can result in spontaneous abortion, repeated pregnancy loss, and stillbirth. Greenberg et al. [30], in a trial from before conception to early pregnancy supplementation with folic acid, they observed that folic acid decreased the risk of NTDs in fetuses. In this study, there was no statistically significant relationship between spontaneous abortion and basophil, neutrophil, lymphocyte and eosinophil counts in UHSA and URSA as compared to the control subjects (Table 2).

**Table 5 Percentage Positivity and Negativity for Antinuclear Antibodies in Blood Group O Subjects**

Parameters	No. of ANAs Positivity	% of ANAs Positivity	No. of ANAs Negativity	% of ANAs Negativity
URSA	3	8.1	34	91.8
UHSA	7	22.5	24	77.4
Control Subjects	3	9.6	28	90.3

**Key:** ANAs = Antinuclear Antibodies

~~This~~ The findings of the present study is are in line with the findings results of Lucca et al. [31], who reported that normal pregnancy is accompanied by leukocytosis caused by physiological stress and that a complex physiological process increases leucocyte counts during pregnancy. According to Donbak et al. [32], the entire human population shares similar ABO and Rhesus blood groups, the frequency and distribution of the blood groups vary among nationalities and races. The distribution of these two blood groups has been repeatedly investigated in various populations all over the world, and their frequencies



exhibited considerable variation in different geographic locations, reflecting the underlying genetic and ethnic diversity of human populations.

Nigeria is a highly populated country comprising of different ethnic groups. As with many other genetic traits, the gene frequency of the ABO and Rh blood group varies significantly within the six geopolitical zones in Nigeria. Hassan et al. [33], reported that, it was revealed that the ABO blood group frequencies were Hassan found in the order O > A > B > AB (52.93%, 22.77%, 20.64% and 3.66%) respectively among Nigeria population. There is no direct relationship between antinuclear antibodies and the ABO blood group, ~~but~~. However, studies ~~has~~ have shown that a high level of antinuclear antibody test is used to measure autoimmune disorder and an autoimmune disorder causes the immune system to attack its own normal cells, tissues and organ by mistake, instead of foreign organisms. As we can see from the result above, since blood type B carries B antigen on its red cells, therefore women with URSA, UHSA & Control participants, with A blood type husbands are likely to have tested positive for antinuclear antibodies because of autoimmune antibodies due to incompatibility. ABO incompatibility happens when a mother's blood type is O, and her baby is A or B. The mother's immune system may react and make antibodies against her baby's red blood cells.

**Table 6 Percentage Positivity and Negativity for Antinuclear Antibodies in Blood Group A & AB Subjects**

Parameters	No of ANAs Positivity		% of ANAs Positivity		No of ANAs Negativity		% of ANAs Negativity	
	A	AB	A	AB	A	AB	A	AB
URSA	0	0	0	0	0	0	0	0
UHSA	0	0	0	0	0	0	0	0
Control Subjects	0	0	0	0	0	0	0	0

**Key:** ANAs = Antinuclear Antibodies

In this study, the percentage positivity for antinuclear antibodies in blood group O subjects (Table 5). Showed 8.1% for those with URSA, 22.5% for UHSA and 9.6% for ANA in control subjects. This agrees with the study of Soni, [34], in which couples who had ~~having~~ O type wives, A or B type husbands, showed an increase in fetal loss. As far as spontaneous abortion is concerned, it is higher in A-A-type husband and O type wife and stillbirths are higher in couples with A-A-type husband and B type wifewives, while Mohanty and Das, [35] discussed the effects of natural selection on four populations of ABO group and found that the incidence of abortion, stillbirth and post-natal mortality are marginally higher among incompatible couple than a compatible couple without revealing significant statistical difference.

**Table 7 Percentage of Hypersegmented Neutrophils in the Study Subjects**

Parameters	No. of Hypersegmented Neutrophils	% Hypersegmented Neutrophils
URSA	18	36
UHSA	15	30
Control Subjects	6	12

**Key:** ANAs = Antinuclear Antibodies

Blood transfusion has been reported to result in high mortality before the discovery of ABO blood groups, ~~;~~ this is because there was no knowledge of the difference in the blood composition among individuals in the human population. ~~Later~~ A later study by Landsteiner made it possible to type/group blood to ABO class (A, B and O) based on the presence or absence of surface antigens on red blood cells (RBC). The last type (AB) was discovered in 1902 by DesCasterllo and Sturli. These germane discoveries lead to reduced mortality due to blood transfusion. The Human ABO type is a classic example of multi-allelism because it

has three alleles (A, B and O), and its phenotype could be O, A, B or AB. The composition for an individual ABO blood type is always based on inheritance of gene on chromosome 9(9q34), which encodes glycosyltransferases that ~~transfersomeoligosaccharides~~ ~~transfersomeoligosaccharides~~ residues to H antigen, resulting in the formation of group A and B antigens, but O individuals lack such activity.

#### 4. CONCLUSION

~~It is therefore concluded that age, ANAs, ABO blood group~~ Therefore, it is concluded that ~~age, ANAs, ABO blood group~~, and monocyte could be responsible for unexplained SA. It is recommended that ANAs, Blood groups and DLC be ~~properly adequately~~ checked in early pregnancy, for this shall further aid the prediction and prevention of spontaneous abortion.

#### CONSENT

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the ~~editorial office/chief editor/editorial board members of this journal~~ ~~is journal's editorial office/chief editor/editorial board members~~. Informed consent of individual participants was sort verbally and with a questionnaire. The importance of the work was pointed out to them and also how it will benefit women generally.

#### ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. The ethical clearance for the study was sourced and obtained from the Ethical Committee of the Federal Medical Center, Yenagoa.

#### REFERENCES

1. Jeve, Y.B. & Davies, W. Evidence-based management of recurrent miscarriages. *Journal Human Reproductive Sciences*, 2014; 7 (3): 12-20.
2. RCOG. The Investigation and Treatment of Couples with Recurrent Miscarriage Guideline. *Royal College of Obstetricians and Gynaecologists*, 2013; 17: 12-31.
3. Oliver-Williams, C. T., Heydon, E. E., Smith, G. C. S. & Wood, A. M. Miscarriage and future maternal cardiovascular disease: A systematic review and meta-analysis. *Heart*. 2013; 99 (22): 1636–44.
4. Goddijn, M. & Leschot, N. J. Genetic aspects of miscarriage. *Baillieres Best Practical Research Clinical Obstetrics and Gynaecology*, 2000; 14: 855-8.
5. Regan, L. & Rai, R. Epidemiology and the Medical Causes of Miscarriage. *Best Practice Research Clinical Obstetrics Gynaecology*, 2000; 14: 839–54.
6. World Health Organization. International Statistical Classification of Diseases and Related Health Problems 10th revision. Geneva, Switzerland. 2006
7. Takeshita, T. Diagnosis & Treatment of Recurrent Miscarriage Associated with Immunologic Disorders: Is paternal lymphocyte immunization a relic of the past? *Journal of Nippon Medical School*, 2004; 71; 308–13.
8. Cervera, R., Font, J., Ramos-Casals, M., García-Carrasco, M., Rosas, J., Morlà, R.M., Muñoz, F.J., Artigues, A., Pallarés, L. Ingelmo, M. Primary

- Sjögren's syndrome in men, clinical and immunological characteristics. *Lupus*, 2000 9 (1): 614-8.
9. Reece, J. & Campbell, N. San Francisco, Pearson/Benjamin-Cummings. *Biology (7th ed.)*. ISBN-13, 978-999. 2005
  10. Janeway C.A Jr, Travers P, Walport M. The immune System in Health & Disease. Immunobiology(5<sup>th</sup> ed.).Newyork:GarlandSciencefromHttp://www.ncbi.nlm.nih.gov/books/NBK 10757 2001.
  11. Capolunghi, F., Rosado, M. M., Sinibaldi, M., Aranburu, A. & Carsetti, R. Why do we need IgM Memory B cells? *Immunology Letters*, 2013; 152 (2): 114–20.
  12. Atanu, Koner., Pallaris, Rajput, Rajat, Dhyani, Nikki Nidhi & Kuljeet Kaur. *Research Journal of Biology (RJB)*, 2013; 1: 36-9.
  13. La Fleur-Brooks, M. Exploring Medical Language, A Student-Directed Approach (7th ed.). St. Louis, Missouri, US. Mosby Elsevier, p. 398 -400. 2008.
  14. Reid ME and Lomas-Francis C. The Blood Group Antigen Facts Book. Second ed. 2004, New York: Elsevier Academic Press.
  15. Marwaha, N., Dhawan, H. & Thakral, B. Severe ABO hemolytic disease of newborn with a positive direct antiglobulin test. *Indian Journal of Pathology Microbiology*, 2009 52: 1-10.
  16. Falah, H. K. Relationship between presence of anticardiolipin antibodies and miscarriage. *Al-Kufa University Journal for Biology*, 2016; 8 (2): 372-83.
  17. Alagoa, E. J. The land and people of Bayelsa State: Central Niger Delta. Onyoma Research Publication, pages 411. 2009
  18. Araoye, M. O. Research Methodology with Statistics for Health and Social Sciences. *Nathadex Publishers, Ilorin*, 2004; 25: 120-5.
  19. Mohammed Hassanzadeh-Nazarabadi, Sahar Shekouhi, & Najmeh Seif. The incidence of spontaneous abortion in mothers with blood group O compared with other blood type. *International Journal of Molecular and Cellular Medicine*, 2012 1(2): 99-104.
  20. Deng, Q., Deng, L., Chen, C., Mengm S. & Pan, Y. Reference Intervals of Hematological Variables for Healthy Antepartum and Postpartum Women. *International Journal of Gynecology and Obstetrics*, 2014; 1: 16–7.
  21. Odotola, A. A., Afolabi, M. O., Jafali, J., Baldeh, I. & Owolabi, O.A. Haematological and biochemical reference values of Gambian infants. *Tropical Medical International Health*, 2014; 9: 275-83.
  22. Sarici, S. U., Alpay, F. & Yesilkaya, E. Hemolytic disease of the newborn due to isoimmunization with anti-E antibodies: a case report. *Turkish Journal of Pediatrics*, 2002; 44: 248–50.
  23. Kutteh, W. H. & Odom L. N. Evaluating Recurrent Pregnancy Loss. *The Role of Autoimmune Screening*, 2012; 10: 5–11.
  24. Rai E, & Wakeland E.k. Genetic predisposition to autoimmunity what have we learned? *Seminars in Immunology*, 2011; 23(2): 67–83.

25. Nyabo Anderson A.M, Wohlfahrt J, Christens P, Olsen J, Melbye M. Maternal age and fetal loss population based register linkage study. *British Medical Journal*, 2000; 320: 1708-12.
26. Segolodi, T. M., Henderson, F. L., Rose, C. E., Turner, K. T. & Zeh, C. Normal Laboratory Reference Intervals among Healthy Adults Screened for a HIV Pre-exposure Prophylaxis. *Clinical Trial in Botswana*, 2014; 9: 930-4.
27. Yang T, Gu Y, Wei X, Liang X, Chen J, Liu Y. Periconceptional folic acid supplementation and Vitamin B12 status in a cohort of Chinese early pregnancy women with the risk of adverse pregnancy outcomes. *Journal of Clinical Biochemical Nutrition*, 2017; 60; 136-42.
28. Thompson M.D, Cole DE, Ray J.G. Vitamin B-12 and neural tube defects: The Canadian experience. *American Journal Clinical Nutrition*, 2009; 89; 697-701.
29. Tamura T, Picciano M.F. Folate and human reproduction. *American Journal of Clinical Nutrition*, 2006; 83: 993-1016.
30. Greenberg J.A, Bell S.J, Guan Y, Yu YH. Folic acid supplementation and pregnancy: More than just neural tube defect prevention. *Review in Obstetrics & Gynecology*, 2011; 4: 52-9.
31. Lucca, L., Rodrigues, F., Jantsch, L. B., Neme, W. S., Gallarreta, F. M. P. Oxidative Profile and  $\delta$ -Aminolevulinatase Dehydratase Activity in Healthy Pregnant Women with Iron Supplementation. *International Journal of Environmental Research in Public Health*, 2016; 13: 463-70.
32. Donbak L, Rencuzogullari E, Topaktas M, Kayrin L. Detection of some blood group (ABO, RH-D) and serum protein (HP, a1-AT, TF) polymorphisms in Antakya Province, Turkey. *TKlin J Med Res*, 2002; 45: 4-9.
33. Hassan, A. Babadoko, A.A., Ahmed, A.J., Isa, H.A., Suleiman A.M. The pattern of distribution of ABO blood groups in North Western Nigeria. *Annals of Nigeria Medicine*, 2005; 1 (2): 17-8
34. Soni, N. & Mukherjee B.M.A Study on Foetal Wastage and ABO Blood Groups Incompatibility among the Gondsof Garriyab and, Chhattisgarh, India. *Anthropologist*, 2009; 11: 229-31.
35. Mohanty, R. & Das, P. K. A Search for Operation of Natural Selection in ABO Blood Groups: Evidences from Four Ethnic Groups of Orissa. *Anthropologist*, 2010; 12: 1-11.